



Key Facts about Swine Influenza (Swine Flu)

What is Swine Influenza?

Swine Influenza (swine flu) is a respiratory disease of pigs caused by type A influenza virus that regularly causes outbreaks of influenza in pigs. Swine flu viruses cause high levels of illness and low death rates in pigs. Swine influenza viruses may circulate among swine throughout the year, but most outbreaks occur during the late fall and winter months similar to influenza outbreaks in humans. The classical swine flu virus (an influenza type A H1N1 virus) was first isolated from a pig in 1930.

Swine Flu in Humans

Can humans catch swine flu?

Swine flu viruses do not normally infect humans. However, sporadic human infections with swine flu have occurred. Most commonly, these cases occur in persons with direct exposure to pigs (e.g. children near pigs at a fair or workers in the swine industry). In addition, there have been documented cases of one person spreading swine flu to others. For example, an outbreak of apparent swine flu infection in pigs in Wisconsin in 1988 resulted in multiple human infections, and, although no community outbreak resulted, there was antibody evidence of virus transmission from the patient to health care workers who had close contact with the patient.

How common is swine flu infection in humans?

In the past, CDC received reports of approximately one human swine influenza virus infection every one to two years in the U.S., but from December 2005 through February 2009, 12 cases of human infection with swine influenza have been reported.

What are the symptoms of swine flu in humans?

The symptoms of swine flu in people are expected to be similar to the symptoms of regular human seasonal influenza and include:

- fever (over 100)
- body aches
- loss of appetite
- coughing
- runny nose
- sore throat
- nausea
- vomiting and diarrhea

Can people catch swine flu from eating pork?

No. Swine influenza viruses are not transmitted by food. You cannot get swine influenza from eating pork or pork products. Eating properly handled and cooked pork and pork products are safe. Cooking pork to an internal temperature of 160°F kills the swine flu virus as it does other bacteria and viruses.

How does swine flu spread?

Influenza viruses can be directly transmitted from pigs to people and from people to pigs. Human infection with flu viruses from pigs are most likely to occur when people are in close proximity to infected pigs, such as in pig barns and livestock exhibits housing pigs at fairs. Human-to-human transmission of swine flu can also occur. This is thought to occur in the same way as seasonal flu occurs in people, which is mainly person-to-person transmission through coughing or sneezing of people infected with the influenza virus. People may become infected by touching something with flu viruses on it and then touching their mouth or nose.

How can human infections with swine influenza be diagnosed?

To diagnose swine influenza A infection, a respiratory specimen would generally need to be collected within the first 4 to 5 days of illness (when an infected person is most likely to be shedding virus). However, some persons, especially children, may shed virus for 10 days or longer. Identification as a swine flu influenza A virus requires sending the specimen to CDC for laboratory testing.

Prevention Steps:

- Avoid close contact with people who may have symptoms or if you have symptoms
- Stay home from work/school when you are sick
- Cover your mouth and nose
- Wash your hands frequently
- Avoid touching your eyes, nose and mouth
- Practice other good health habits

What to do if you have symptoms:

- Call your doctor to get guidance on what he/she wants you to do.

What medications are available to treat swine flu infections in humans?

There are four different antiviral drugs that are licensed for use in the US for the treatment of influenza: amantadine, rimantadine, oseltamivir and zanamivir. While most swine influenza viruses have been susceptible to all four drugs, the most recent swine influenza viruses isolated from humans are resistant to amantadine and rimantadine. At this time, CDC recommends the use of oseltamivir or zanamivir for the treatment and/or prevention of infection with swine influenza viruses.

FOR MORE INFORMATION, PLEASE CONTACT BLACK HEALTH COALITION OF WISCONSIN AT 414-933-0064 EXT. 207 OR THE MILWAUKEE HEALTH DEPARTMENT'S HOTLINE AT 414-286-3616

(Source: CDC, Questions & Answers: Key Facts about Swine Influenza (Swine Flu))